

2019.12.11

福岡市西区・今津干潟のヘラサギ、ミサゴ、ハヤブサ



ヘラサギ(籠鷺)
トキ科 L=86cm



カワウ(河鵜)
ウ科 L=82cm

ヘラサギ(篋鷺)
トキ科 L=86cm







ミサゴ(鶺鴒) ワシタカ科 L=メス64cm



ミサゴ(鶺鴒) ワシタカ科 L=メス64cm



ミサゴ(鶺鴒)
ワシタカ科 L=メス64cm



ヘラサギ(篋鷺)
トキ科 L=86cm



クロツラヘラサギ(黒面籠鷺)
トキ科 L=77cm



クロツラヘラサギ(黒面篋鷺)
トキ科 L=77cm



ミサゴ(鵟) ワシタカ科 L=メス64cm



ミサゴ(鵟) ワシタカ科 L=メス64cm



ミサゴ(鵟) ワシタカ科 L=メス64cm



ズグロカモメ(頭黒鷗)
カモメ科 L=32cm



ホウロクシギ(焙烙鷓)
シギ科 L=63cm



ミサゴ(鷺) ワシタカ科 L=メス64cm




クロツラヘラサギ(黒面籠鷺)
トキ科 L=77cm



ヘラサギ(籠鷺)
トキ科 L=86cm





コサギ(小鷺)
サギ科 L=61cm

クロツラヘラサギ(黒面篋鷺)
トキ科 L=77cm



クロツラヘラサギ(黒面篋鷺)
トキ科 L=77cm



クロツラヘラサギ(黒面篋鷺) トキ科 L=77cm



イソシギ(磯鶺)
シギ科 L=20cm



ムクドリ(椋鳥) ムクドリ科 L=24cm



ハヤブサ(隼)
ハヤブサ科 L=メス49cm


A photograph of two white egrets standing in shallow water. The larger egret on the left is facing right, and the smaller egret on the right is facing left. The water is dark and rippled, and the ground they are standing on is dark and muddy. The background is a soft, hazy sky.

ダイサギ(大鷺)
サギ科 L=90cm


コサギ(小鷺)
サギ科 L=61cm

A Hamashigi bird, also known as a beach plover, is shown standing in shallow, rippling water. The bird has a long, dark, straight bill and a body with mottled brown and white feathers. The water is clear, and the background shows gentle waves. The lighting suggests a late afternoon or early morning setting.

ハマシギ(浜鷗) シギ科 L=21cm



イノシギ(磯鶺)
シギ科 L=20cm



イノシギ(磯鷗)
シギ科 L=20cm

END