


2020.04.20 福岡市西区・今津干潟の
ハマシギ、キアシシギ、ソリハシシギ、クロツラヘラサギ



A Hamashigi bird (濱鷗) is shown in profile, facing left. It has a long, dark, straight beak and a body covered in intricate brown and white mottled patterns. The bird is standing on a dark, wet rock covered in green seaweed. The background is a shallow, rippling body of water with more seaweed scattered throughout.


ハマシギ(濱鷗)
シギ科 L=21cm



ヒドリガモ (緋鳥鴨)
ガンカモ科 L=49cm



ヒドリガモ(緋鳥鴨)メス
ガンカモ科 L=49cm



キアシシギ (黄足鷗)
シギ科 L=25cm



キアシシギ (黄足鷗)
シギ科 L=25cm



アオアシシギ (蒼脚鷗)
シギ科 L=35cm

ダイゼン (大膳)
チドリ科 L=29cm



ツルシギ (鶴鷗)
シギ科 L=30cm



ダイゼン (大膳)
チドリ科 L=29cm



ソリハシシギ (反嘴鷸)
シギ科 L=23cm

クロツラヘラサギ (黒面篋鷺)
トキ科 L=77cm



クロツラヘラサギ (黒面篋鷺)
トキ科 L=77cm



クロツラヘラサギ (黒面篋鷺)
トキ科 L=77cm



A group of shorebirds, including a grey-brown bird and several brown and white speckled birds, are wading in shallow water. They are surrounded by large, dark, moss-covered oyster shells. The water is rippled, and the birds are engaged in foraging or resting. The overall scene is a natural coastal habitat.

キアシシギ (黄足鷸)
シギ科 L=25cm

ハマシギ (浜鷸)
シギ科 L=21cm

毘沙門山（びしゃもんやま）177m

13:04

浜崎山 (はまさきやま) 97m

13:04

柑子岳（こうしだけ）255m

13:08

END